

Evolutionary route to diploidy and sex

Erkan Tüzel^{1,2}, Volkan Sevim¹ and Ayşe Erzan^{1,3}

¹ *Department of Physics, Faculty of Sciences and Letters
Istanbul Technical University, Maslak 80626, Istanbul, Turkey*

² *Department of Physics, Faculty of Sciences and Letters
Işık University, Maslak, 80670, Istanbul, Turkey*

³ *Gürsey Institute, P.O.B. 6, Çengelköy, 81220 Istanbul, Turkey*

Using a bit-string model of evolution, we find a successful route to diploidy and sex in simple organisms. Allowing the sexually reproducing diploid individuals to also perform mitosis, as they do in a haploid-diploid cycle, leads to the complete takeover of the population by sexual diploids. This mechanism is so robust, that even the accidental conversion and pairing of only two diploids give rise to a sexual population.

Keywords: Evolution, sexual reproduction, haploid-diploid cycle.

I. INTRODUCTION

The evolution of sex through Darwinian selection in spite of the seeming odds with respect to asexual reproduction through cloning, is a still unresolved problem [1, 2, 3], which has attracted a great deal of recent activity [4, 5]. We would like to present here results from a very simple model based on unicellular organisms where sex seems to have first emerged, without any complicating factors such as the cost of male-female differentiation, parental care, mating preferences, maternity periods and so on, which we assume only to have arisen in higher organisms. These factors may be responsible for the differentiation, modification or even at times reversal (as in meiotic parthenogenesis [2], p.234) of the patterns that may have become accidentally established in the remote evolutionary past. Unicellular organisms are not only simpler to model, but this choice also imposes more stringent conditions on the establishment of a sexual population, since it severely limits the number of viable offspring that a pair of parent cells can have. By confining ourselves to “worst case scenarios” we hope to be able to obtain plausible lower bounds to the feasibility of sex as a reproductive mechanism.

Sexual reproduction as we know it today in higher plants and animals can be regarded as a haploid-diploid cycle with a highly abbreviated haploid phase, where the (haploid) gametes typically do not perform mitosis. [1, 2] These haploids are not viable unless two of them fuse and once more make a diploid cell, which then multiplies by mitosis and eventually forms the mature individual.

The haploid diploid cycle (HDC) where, under unfavorable conditions the population becomes diploid, is found in many species [6, 7], and was the motivation behind the hypothesis of Jan, Stauffer and Moseley [8] (JSM) who proposed that diploidy and sex may have first arisen as a way to escape death, when a simple, unicellular individual is threatened by too many deleterious mutations. The fitness is taken to be a step function of the number of mutations, also setting the threshold for conversion to diploidy and sex. Here, the number of deleterious mutations can be read as the distance from

the ideal case, called the wildtype, which can be altered as a result of a change in the environment, so that the individual is not as well adapted as before the change.

In previous work [9, 10] we showed that the JSM hypothesis indeed leads to a steady source of diploid sexuals, given a population of haploids which multiply by mitosis. This leads to a steady state distribution of coexisting haploids and diploids in a constant population. The premises adopted in [10] were very restrictive. Two sexual “parent” cells were allowed to have only one offspring, after which they died. In case one considers greater number of offspring, by allowing a greater number of viable gametes to be formed, as is indeed possible in many unicellular organisms [6], one finds that the diploid, sexual population completely takes over.

In the same paper [10], we have also considered the situation where conversion to sex occurred with a constant probability σ (which we varied between 0 and 1), independently of the fitness of the individual. This strategy also gave rise to a sexual population making up a small but macroscopic fraction of the total, so that the JSM threshold mechanism for conversion to sex did not prove to be necessary, although it was more successful. On the other hand the JSM hypothesis has the additional attraction of providing a mechanism which could trigger the fusion of two haploid cells to form a diploid: it is known [1], p.149, that extensive genetic damage can lead to gene repair via genetic transmission between two haploid cells; the fusion of two haploids could be seen as an extreme form of such behaviour. An alternative means of forming a diploid from a haploid cell is via endomitosis [2], p.230, as in the first step in meiotic parthenogenesis. Endomitosis is the process whereby the genetic material in a cell is duplicated without subsequent cell division as in normal mitosis. Again, it is not implausible that this process originated as a result of grave genetic damage which precluded the successful completion of mitosis. [1], p.149.

In the present paper we show that, if the diploid cells, once formed, are also allowed to multiply by mitosis, as indeed they do in a haploid-diploid cycle, the whole population is taken over by diploid cells, who perform facultative sex if they are once more threatened by extinction

due to too many deleterious mutations. Moreover we show that even an episodic conversion to sex, involving as few as only two individuals who survive to mate, leads to a steady state made up solely of sexual types.

In the next section we briefly describe our algorithms and report the simulation results. In section 3, we provide a discussion and some pointers for future research.

II. BIT-STRING MODEL FOR THE CONVERSION TO SEX - ALGORITHMS AND SIMULATION RESULTS

The type of model we consider here is the same as in Ref. [10]. Below and in Section A, we recall the definitions. In Section B and C we introduce new rules to test the autonomous viability of sexual populations.

Each haploid one-celled organism consists, for our purposes, of a 15-bit string of “0”s and “1”s, representing the genetic code in a 16-bit computational word. [11, 12, 13] We use the bit defining the “sign”, to specify whether the individual is asexual (+) or sexual (-). A mutation consists of flipping a randomly chosen bit except the sign bit, with a constant probability, Γ , for each individual per generation. Since the genetic difference between individuals of the same species is typically less than 10% [1], p.52, this rather short string for the genetic code may be considered as a coarse grained model for the complete genome of the individual, which we divide up into different zones, retaining a “0” where there are no mutations, and flipping the bit to “1” if there are one or more mutations in this zone. The wild type is a string of all “0”s. Therefore, at each locus, a “1” corresponds to a deleterious mutation (which we will call “mutation,” for short, where this is not liable to lead to any confusion.) Diploid organisms have two bit strings, which are allowed to be different. The number of deleterious mutations m is simply the number of “1”s for a haploid individual. For a diploid, the number of “expressed” deleterious mutations is the number of loci at which both homologous alleles are set to “1,” i.e., we assume that deleterious mutations are recessive.

The total population is fixed at $N = 10^3$, and we have chosen $\Gamma = 1/N$. This corresponds to a mutation rate per allele per generation of $\sim 6 \times 10^{-5}$, which is comparable to the typical mutation rates encountered in eukaryotes [2], p.64.

The probability of survival (or fitness) as a function of m is given by a Fermi-like distribution [14], $P(m)$,

$$P(m) = \frac{1}{\exp[\beta(m - \mu)] + 1} \quad (1)$$

For large β (or “low temperatures,” in the language of statistical mechanics), $P(m)$ behaves like a step function. Individuals with $m > \mu$ die, those with $m < \mu$ survive, and those with $m = \mu$ survive with a probability of 1/2. In the simulations we confined ourselves to low temperatures ($\beta = 10$). We chose $\mu = 4$ which just allows us

enough variability without leading to totally unrealistic mutational loads.

We start with a set of N asexual (haploid) wildtypes. In each generation about ΓN individuals suffer mutations; they are killed off or retained according to the fitness function (1), and the population restored to N by once cloning as many randomly chosen survivors as necessary. The population of haploid asexuals settles down to a minimally stable [15] steady state distribution (see Table I) as shown in Fig. 1, independently of the value of Γ , for $\Gamma \geq 1/N$. [9, 10]

For comparison, we have also performed simulations on a diploid population reproducing asexually, according to the same rules as stated in the previous paragraph. We have found that the diploid population reaches a steady state with an m distribution peaked at $m = 2$, rather than $m = 3$ as found for haploids. (See Fig. 1 and Table I).

A. Conversion to sex

Let us briefly summarise our algorithm for conversion to sex, which we also used in our previous study (see Model A, Ref. [10]).

We choose the steady state of the haploid population as our initial state from which to start the conversion to sex, as that would be a most likely “natural state” encountered at this stage. Once the haploid population reaches a steady state, we allow those individuals that are threatened by extinction to convert to diploidy and sex, by implementing the following rule at each generation: If an asexual individual with $m = 4$ has survived, it is converted to an active sexual, by deterministically and irreversibly switching its sign bit to (-). It performs endomitosis and becomes diploid. If there are already sexual, diploid organisms in the population, they will also be made active if m , the number of their expressed mutations, is ≥ 4 (otherwise they do not participate in the reproduction cycle; hence the conversion to sex may be termed non-hereditary). Finally, all the active sexual organisms pair randomly and engage in sexual reproduction, where they each contribute one gamete (formed via one step meiosis) towards a single diploid sexual offspring. If we denote the genotypes of parents as $\{Aa\}$ and $\{Bb\}$ respectively, then the genotype of the offspring is either $\{AB\}$, $\{Ab\}$, $\{aB\}$ or $\{ab\}$. No crossover occurs during this one step meiosis. If there is only one active sexual at a certain time step then it must wait subsequent generations until it either finds a partner or it dies. We keep the population constant [16] by cloning randomly selected *haploid* individuals to make up the deficit.

(The consequences of the conversion to hereditary and obligatory sex can be found in Ref. [10].)

B. Mitotic Diploid Sexuals Win over Haploids

In the previous subsection we described a scenario where too big a mutation load meant conversion to diploidy and sex, if the individual survived. It should be realised that in this scheme the haploids, who multiply by cloning, provide a steady source for the diploid sexuals, whose numbers are halved every time they mate. Now we want to test whether the diploid, sexual population can survive autonomously, if the sexuals are also allowed to perform mitosis themselves.

Once a steady state with coexisting haploid and diploid populations is reached via the algorithm described above, we switch off the conversion of haploid organisms to diploidy and sex. The diploid individuals mate when they face extinction due to too many mutations. We now keep the total population constant by making up for the deficit in the population at each step by cloning randomly selected individuals, *regardless of whether they are sexual or asexual*. In colonies undergoing a haploid-diploid cycle, it is quite frequently the case [6] that the diploid phase of the cycle also involves multiplication by mitosis.

The result is that the diploid, sexual individuals completely win over the population. The haploids which now can not compete with the sexuals, become extinct. Here we find that the haploid phase of the HDC becomes abbreviated to the point where haploids appear only as gametes which do not perform mitosis. This is exactly the situation in highly evolved sexual organisms.




FIG. 1: The steady state distribution of pure haploids with respect to the number deleterious mutations m , compared with the distribution of asexual diploids over the number of *expressed* mutations m . See text. The histograms are normalized to unity.

C. Episodic conversion to diploidy and sex leads to sexual steady state

One can also think of a scenario in which conversion to sex takes place accidentally over a short period of time after the asexual population settles down to a steady state. In fact this is probably the most realistic situation, given the random nature of the mutations.

The way we actually implemented this in the computer code was by deterministically switching the sign bits of the first two asexual individuals to survive with $m = 4$, and then turning off the possibility of further conversion. These then form two sexual individuals by endomitosis, and if the first survives long enough so that it can mate with the second, will give rise to one sexual offspring.

The rest of the rules are as explained before; in each generation we clone randomly chosen individuals to make up the deficit in the population, regardless of whether they are asexuals or sexuals. We allow the diploid individuals to mate when they face extinction in the course of their lives.

Suprisingly, the sexuals capture the population in 95 percent of the performed runs. (In the rest, the single diploid, which is still at the threshold of extinction with an m value of 4, may not survive until a partner arrives.)

D. Relative Fitness

The steady state m distributions for the sexual populations treated in Section II.B and C are much better than the haploid distribution. However, they are identical with the distribution for asexual diploids. (see Table I and Fig. 1) This means that as soon as diploidy is achieved, it is so successful in screening the effects of deleterious mutations that once diploid, the organism practices sex very infrequently, making these three distributions identical.

The diploid m -distribution in Fig. 1 should be contrasted with the results of Ref. [10] (Model A, non-hereditary and hereditary; also see Subsection A, above) where cloning of the sexuals is not allowed, but there is a steady influx of new sexuals from the haploids (see Table I). In the non-hereditary model, the m distribution has shifted nearer the wildtype than in the purely diploid case (although the peak is again at $m = 2$); in the hereditary model, the peak has shifted to $m = 1$. Thus, the frequency with which sex is practiced has a salutary effect on the distribution of expressed deleterious mutations in the population.

On the other hand it should be noted that, with the Fermi-like fitness function (Eq.(1)) for large β , to which we have confined ourselves, the survival probability is $\sim$ constant for $m < \mu$, and therefore does not discriminate between steady state populations that differ solely in the shape of their m distributions for $m < \mu$. A better measure might be the deathrate, namely, the average number of individuals which are eliminated at each step, under

TABLE I: The distribution of expressed deleterious mutations and death rates for the models discussed in Section II. The m -distributions are percentages ($\pm 1\%$) computed over each type if coexisting; the death rates are per capita per generation. (* see Ref. [10])

		$m = 0$	$m = 1$	$m = 2$	$m = 3$	$m = 4$	death rate
Pure Haploid		1	10	36	53	0	$O(10^{-4})$
Pure Diploid		7	27	40	26	0	$O(10^{-5})$
Diploid Sexuals	Mitotic	7	27	40	26	0	$O(10^{-5})$
Diploid Sexuals	Episodic	7	27	40	26	0	$O(10^{-5})$
Model A - non-hereditary*	Asex	1	11	37	51	0	$O(10^{-4})$
Model A - non-hereditary*	Sex	11	33	36	20	0	$O(10^{-4})$
Model A - hereditary*	Asex	1	11	36	52	0	$O(10^{-4})$
Model A - hereditary*	Sex	10	50	40	0	0	$O(10^{-2})$

the criterion given in Eq.(1).

Our findings, which are rather revealing, are summarized in Table I. Since the fluctuations were very large, we have only reported order of magnitude results for the death rates. Pure diploids have the lowest death rate, an order of magnitude smaller than pure haploids. Mitotic (including the “episodic” case) sexuals are very close to diploids. For non-hereditary sexuals the rates are comparable to asexuals, but hereditary sexuals have a death rate that is two orders of magnitude larger than asexuals. We infer from this that the success of hereditary sexuals in lowering their mutation load is due to the much greater rate at which they can select-out highly mutated genes through death.

Thus it is seen that the clear advantage of sex over asex (haploid or diploid) in our model can only become manifest with a sufficiently high frequency of sexual reproduction, which is driven by the mutation rate Γ , and moreover in situations which can discriminate between the different adaptabilities (smaller typical m) of the different modes of reproduction, such as time varying environments. [17] This is similar to the finding of Pekalski [18], who has considered environments (wildtypes) that vary over time, and found that the benefits of sexual reproduction are enhanced by higher birthrates. For lower birthrates, meiotic parthenogenesis, which can be compared to diploid asex, has a slight advantage over sex, a difference from our results which can be ascribed to the exponential fitness function he uses.

III. DISCUSSION

This paper is a culmination of a series of studies where we have considered very stringent rules for the survival and propagation of diploid, sexual individuals in competition with haploids. Here we have finally been able to show that a pair of simple, unicellular organisms who have accidentally converted to diploidy, and which subsequently engage in sexual reproduction, begetting one sexual offspring, can give rise to a population of sexual types which totally take over a finite population, provided they are also allowed to multiply by mitosis, on an

equal footing with the haploids in the population. They engage in sex when the going gets tough, that is, when the number of their expressed deleterious mutations exceeds a certain number. This success seems to vindicate the hypothesis of JSM [8], that sex could have been a mechanism of last resort when simple organisms were faced with extinction.

It is interesting to pose the question of what happens if we do not allow the diploids to clone themselves, but, on the other hand, allow the gametes, under special conditions, to enter a haploid phase where they multiply by mitosis. Eventually these haploids will be allowed once more to fuse and give rise to diploids. Within the present scheme, since the haploids convert to diploidy only at the threshold, $m = 4$, the gametes of the diploid individuals have $m \geq 4$, i.e., they are not viable. Therefore the extension of the haploid phase calls for a modification of the rules in such a way as to allow the gametes to survive nevertheless, e.g., by forming “spores.”

One way the parameters of the present model could be modified is to take the threshold for survival to be different (greater than) the threshold for the conversion to (or practice of) sex. Then, if a haploid gamete were to be considered as a member of the haploid population, and subjected to the same rules for selection and reproduction, it would seem as if one would get haploid phases of arbitrary duration. However, upon closer inspection we see that this is not true: Since the gamete comes from a diploid organism which engages in sex due to an excess mutational load, it already has deleterious mutations in excess of (or at least equal to) the sex threshold. (For the diploid organism to have m expressed mutations, both of its genetic strings must have at least that many). So, according to the rules which we have adopted, the abbreviated haploid cycle is automatically selected, since within this scheme the gamete would immediately seek a mate. The lengthening of the haplophase would call for further evolution, e.g., such mutations as would lead to the enhancement of the fitness of the haploid gametes with m in excess of the sexual threshold (say by the formation of spores, to survive in an environment which has become disfavorable [3,4]). Since this will provide a mechanism for the diploids once more to make up for the reduction

in their numbers due to the single-offspring mating rule, we expect this HDC to win out over the haploid, asexual population.

To understand the seemingly contradictory of results from similar models investigating the phenomenon of the selection of sex as the dominant mode of reproduction, it is of interest to compare our model with that of Redfield [16] and Cui et al., [19]. The former employs a more complex system in which there is sexual differentiation between males and females, with the additional feature of a much greater mutation rate for males than for females, eventually making meiotic parthenogenesis safer than sex. Redfield has used three different “selection functions” (survival probabilities), falling off exponentially (no “epistasis,” or correlation between the effects of successive mutations) quadratically (some positive epistasis) or as a step function (extreme positive epistasis) with m [20]. The fact that the step-function (“truncation”) survival probability alone leads to a much better fitness resulting from sex, tells us that the form of the survival probability which is adopted determines the outcome very strongly. The step function favors sex because it does not punish small deviations from the wildtype, thereby allowing genetic diversity, while it severely penalizes large deviations, helping eliminate highly mutated individuals.

This insight enables us to understand the results of Cui et al. who have considered a diploid population reproducing either asexually or sexually, depending upon a fixed probability. There is a constant mutation rate. The survival probability is chosen to be of the “independent mutation,” i.e., exponential type. These authors find that the diploid population accumulates such a large number of deleterious mutations (which however do not get expressed), that sexual reproduction, i.e., the random pairing of gametes from different parents, results in a disastrous reduction in the fitness of the offspring. Moreover, sexually reproducing diploid populations are susceptible to invasion by asexually reproducing ones, while the converse is not true, i.e., asexually reproducing diploid populations are not invaded by sexually reproducing ones. These results change drastically in favor of sex when they introduce cell senescence, which means a cell can clone itself only a certain number of times before it stops dividing and dies, unless it engages in sexual reproduction, which resets the senescence clock.

We would like to argue that in the model of Cui et al., introducing cell senescence is very much like turning on a “truncation like” survival function, which, as in our model, also controls the switch to sexual reproduction. In the presence of a constant mutation rate, the number of times a cell has cloned is another way to measure its (mean) mutation load, which, in an asexual haploid

population of course gets directly expressed, whereas for diploid individuals hides behind the dominant unmutated alleles. Introducing a cell senescence threshold, beyond which the individual is either killed off or has to engage in sexual reproduction, is therefore equivalent, in an average way, to the JSM criterion, applied to a diploid population. Seen in this way, it is very gratifying that the Cui et al. model in fact corroborates our findings regarding the greater fitness of diploid populations that frequently engage in sex, given a step-function like survival probability.

Very recently, two variants of the Redfield model, which assign “harsher selection” to males (rather than relatively higher mutation rates) have been considered [21, 22]. These studies both find that the mutational load on the whole population decreases as a result, and that the relative fitness of sexual females is increased by more than enough to compensate for the twofold cost of sex in anisogamous populations, at least for extremely high values of the average mutation rate. Agarwal finds that this can happen even in the absence of synergistic epistasis, in his model. Wilke et. al [23] have meanwhile found that high mutation rates select for low replication rates and flat regions of the fitness surface.

It should be noted that the Fermi-type fitness function (1) adopted here[8, 9, 10, 14] extrapolates, for finite temperatures (smaller values of β) between highly synergistic (step function) and independent (exponential) survival probabilities[24] as a function of m . A recent study by Peck and Waxman indicates [25] that competition for limited resources can lead to synergy between successive mutations, leading to truncation, or step function-like survival probabilities, which they also find favors sexual populations.

In this paper we have presented numerical results for a model involving unicellular organisms, which may shed light on how sex and diploidy emerged and established a foothold in the protozoon world. Many different mechanisms have so far been proposed via which sex may prove advantageous or otherwise, in more highly evolved organisms. We believe that care should be taken while proposing any single mechanism, such as an adaptation to resist infestation by parasites [26, 27], for the preference of sex by organisms that range from the unicellular eukaryotes to trees or human beings [28]. More complex organisms may have elaborated much more complex survival mechanisms and behavioral patterns, which stabilize or destabilize already evolved traits.

Acknowledgements

We thank Aslıhan Tolun and Canan Tamerler Behar for many useful discussions. One of us (A.E.) gratefully acknowledges partial support from the Turkish Academy of Science.

[1] J. Maynard Smith and E. Szathmary, *The Major Transitions in Evolution* (Oxford University Press, Oxford,

1995).

- [2] J. Maynard Smith, *Evolutionary Genetics* (Oxford University Press, Oxford 1999) p.225ff.
- [3] S.C. Stearns, *The Evolution of Sex and its Consequences* (Birkhäuser, Basel, 1987).
- [4] P. Hines and E. Culotta, *Science* **281**, 1979 (1998), special issue; see references therein.
- [5] B. Drossel, *Adv. Phys.* **50**, 209 (2001).
- [6] L.M. Prescott, J.P. Harley, D.A. Klein, *Microbiology* (McGraw Hill, NY 1999) p. 533.
- [7] E.P. Solomon, L.R. Berg, D. Martin, *Biology* (Saunders College Publ., USA 1999) p. 31-33.
- [8] N. Jan, D. Stauffer and L. Moseley, *Theory in Biosciences* **119**, 166 (2000).
- [9] B. Örcal, E. Tüzel, V. Sevim, N. Jan and A. Erzan, *Int. J. Mod. Phys. C*, **11**, 973(2000). See references therein.
- [10] E. Tüzel, V. Sevim, A. Erzan, cond-mat/0101426.
- [11] D. Stauffer, *Physica A*, **273**, 132 (1999).
- [12] T.J. Penna, *J. Stat. Phys.* **78**, 1629 (1995).
- [13] H.L. Muller, *Mutat. Res.* **1**, 2 (1964).
- [14] J. Thoms, P. Donahue, N. Jan, *J. Physique I* **5**, 935 (1995).
- [15] C. Chisholm, N. Jan, P. Gibbs, and A. Erzan, *Int. J. Mod. Phys. C* **11**, 1257(2000).
- [16] R.J. Redfield, *Nature (London)*, **369**, 145 (1994).
- [17] V. Sevim and A. Erzan, to be published.
- [18] A. Pekalski, *Eur. Phys. J. B* **13**, 791 (2000).
- [19] Y. Cui, R.S.Chen, and W.H Wong, *Proc. Natl. Acad. Sci.* **97**, 3330 (2000).
- [20] A. Kondrashov, *Nature* **336**, 435 (1988).
- [21] S. Siller, *Nature* **411**, 689 (2001).
- [22] A.F. Agarwal **411**, 692 (2001).
- [23] C.O. Wilke, J.L. Wang, C. Ofria, R.E. Lenski, and C. Adami, *Nature* **412**, 331 (2001).
- [24] S.F. Elena and R.E. Lenski, *Nature* **390**, 395 (1997).
- [25] J.R. Peck and D. Waxman, *Nature* **406**, 399 (2000).
- [26] R. S. Howard and C.M. Lively, *Nature* **367**, 554 (1994).
- [27] J.S. SaMartins, *Phys. Rev. E* **61**, 2212 (2000).
- [28] W.D. Hamilton, R. Axelrod and R. Tanese, *Proc. Natl. Acad. Sci.* **87**, 3566 (1990).